

## CS 30: Discrete Math in CS (Winter 2020): Lecture 29

Date: 2nd March, 2020 (Monday)

Topic: Infinite Sets: Countability

*Disclaimer: These notes have not gone through scrutiny and in all probability contain errors.*

*Please discuss in Piazza/email errors to deeparnab@dartmouth.edu*

---

1. **Examples of some infinite sets.** A set  $S$  is an infinite set if  $|S| = \infty$ . What does that mean? Well, it means that for *any* natural number  $N$ , one can find  $> N$  distinct elements of  $S$ . Here are some examples of infinite sets we will see the next two lectures.

(a) *The Naturals.*  $\mathbb{N} = \{1, 2, 3, \dots\}$

(b) *The Integers.*  $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\} = \{x : x \in \mathbb{N}\} \cup \{-x : x \in \mathbb{N}\} \cup \{0\}$

(c) *The Rationals.*  $\mathbb{Q} = \{\frac{p}{q} : p \in \mathbb{Z}, q \in \mathbb{Z} \setminus \{0\}\}$

(d) *The Reals.*  $\mathbb{R}$ . What are the reals? That is a deep question and forms the first few lectures of an Analysis course. For us, we will go with

$$\mathbb{R} = \left\{ \sum_{i=0}^{\infty} \frac{a_i}{10^i}, \quad a_0 \in \mathbb{Z}, \quad 0 \leq a_i \leq 9, \quad \forall i \geq 1 \right\}$$

The numbers  $a_1, a_2, \dots$  form the *decimal notation* of the number denoted as the summation.

(e) *Python Programs.*  $\mathcal{P}$ . The set of all possible Python programs.

(f) *The Strings.*  $\Sigma^*$ . The set of all strings formed by using letters from a *finite set*  $\Sigma$ .

(g) *Boolean Functions.*  $\mathcal{F}$ . The set of all functions which assign each natural number a value either 0 or 1.

$$\mathcal{F} := \{f : \mathbb{N} \rightarrow \{0, 1\}\}$$

For instance the `isPrime( $n$ )` function is an element of  $\mathcal{F}$ .

There are two main points to this and the next lecture.

- There are many kinds of infinities (we will see two).
- The cardinalities of the set of Boolean functions, and the set of Python programs are *different!* Thus, there *must* exist functions which have no programs.

And then, lastly, we will see an *explicit* problem which cannot have any algorithm.

2. **Recall.** A function  $f : A \rightarrow B$  is an *injection* if for any two distinct  $a_1 \neq a_2$  in  $A$ , we have  $f(a_1) \neq f(a_2)$ .

3. **Countable Sets.** A set  $S$  is called **countable** if there exists an injection  $f : S \rightarrow \mathbb{N}$ .

It is called so because the elements of  $S$  can be ordered and counted one at a time (although the counting may never finish). More precisely, using  $f$  we can devise an ordering of the elements of  $S$  and an algorithm which for any natural number  $k$  gives the  $k$ th number in the ordering. The following code prints this sequence.

```

1: for  $n = 1, 2, 3, \dots$  do:  $\triangleright n \in \mathbb{N}$ 
2:   if there exists some  $a \in S$  such that  $f(a) = n$  then:  $\triangleright$  i.e.  $f^{-1}(n) \in S$ 
3:     Print  $a$ 

```

Different injective functions can lead to different orderings.

#### 4. Examples of Countable Sets.

- *Finite sets* are trivially countable. If a set  $S$  is finite and  $|S| = k$ , then the elements of  $S$  can be renamed as  $\{e_1, e_2, \dots, e_k\}$ . The injective function  $f(e_i) = i$  implies  $S$  is countable.
- $\mathbb{N}$  is countable by definition. But there are many more interesting examples.
- *Set of Integers.* The set  $\mathbb{Z}$  is countable. To see this, consider the following function  $f : \mathbb{Z} \rightarrow \mathbb{N}$ . If  $x > 0$ , then  $f(x) = 2x$ . If  $x \leq 0$ , then  $f(x) = 2(-x) + 1$ . Note that the co-domain of this function is indeed the natural numbers.

For instance,  $f(2) = 4$ ,  $f(-2) = 5$ , and  $f(0) = 1$ .

**Claim 1.** The function  $f : \mathbb{Z} \rightarrow \mathbb{N}$  defined above is injective.

*Proof.* To see this is injective, we need to show  $f(x) \neq f(y)$  for two integers  $x \neq y$ . We may assume, without loss of generality,  $x < y$ . If both  $x$  and  $y$  are positive, then  $f(x) = 2x < 2y = f(y)$ . Similarly, if both are non-negative, then we get  $f(x) = -2x + 1 > -2y + 1 = f(y)$ . The only other case is  $x$  is non-negative and  $y$  is positive. In this case,  $f(x)$  is odd while  $f(y)$  is even.  $\square$

If we use the above algorithm to figure out the ordering of  $\mathbb{Z}$ , we get:

$$(0, 1, -1, 2, -2, 3, -3, 4, -4, \dots)$$

#### 5. Some operations that preserve countability.

**Theorem 1.** If  $S$  is countable, and  $T \subseteq S$ , then  $T$  is countable.

*Proof.* If  $f : S \rightarrow \mathbb{N}$  is an injection, then the restriction of  $f$  to  $T$ , that is,  $g : T \rightarrow \mathbb{N}$  defined as  $g(t) = f(t)$  is also an injection.  $\square$

**Theorem 2.** If  $S$  is countable and  $T$  is countable and  $S \cap T = \emptyset$ , then  $S \cup T$  is countable.

*Proof.* We can use the trick for showing integers are countable.

Let  $f : S \rightarrow \mathbb{N}$  be the injective function and  $g : T \rightarrow \mathbb{N}$  be the injective function which show they are countable. We now define a function  $h : S \cup T \rightarrow \mathbb{N}$  which is injective. Indeed,

$$h(x) = \begin{cases} 2f(x) & \text{if } x \in S. \\ 2g(x) + 1 & \text{if } x \in T. \end{cases}$$

To prove this is an injective function, take any two  $a \neq b$  in  $S \cup T$ . Either both are in  $S$ , in which case  $h(a) = 2f(a) \neq 2f(b) = h(b)$  where  $f(a) \neq f(b)$  for  $f$  is an injection. Similarly, if both are in  $T$ , then  $h(a) \neq h(b)$ . If one is in  $S$  and the other is in  $T$ , then  $h(a)$  (if  $a \in S$ ) is even while  $h(b)$  is odd. Thus,  $h(a) \neq h(b)$  here as well.  $\square$

**Theorem 3.** If there is a function  $g : A \rightarrow B$  which is an injection, and the set  $B$  is countable, then the set  $A$  is countable.

*Proof.* Since  $B$  is countable, there is an injective function  $f : B \rightarrow \mathbb{N}$ . We claim that the function  $(f \circ g)$  is an injective function from  $A$  to  $\mathbb{N}$ . Indeed, if  $a \neq a'$ , then  $g(a) \neq g(a')$ . Let  $b = g(a)$  and  $b' = g(a')$ . We get  $(f \circ g)(a) = f(b)$  and  $(f \circ g)(a') = f(b')$ . Since  $b \neq b'$ , we get  $(f \circ g)(a) \neq (f \circ g)(a')$ .

□

6. **The Set of Rationals is Countable.** This may be a surprise since the set of rationals are dense, that is, between any two rational numbers, there is a rational number. Nevertheless, they are countable.


To show this, we need to construct an injection  $g : \mathbb{Q} \rightarrow \mathbb{N}$ . For now, we only show an injection of  $g : \mathbb{Q}_+ \rightarrow \mathbb{N}$  where  $\mathbb{Q}_+$  are all the positive rationals; we leave the extension to the full set of rationals as an exercise.


This can be defined as follows: given any positive rational number  $z = p/q$  in the *reduced form* (that is,  $\gcd(p, q) = 1$ ), define

$$z = p/q \quad g : z \mapsto 2^p 3^q$$

Clearly, the functions maps a positive rational number to a positive integer.

We claim that the above function  $g : \mathbb{Q}_+ \rightarrow \mathbb{N}$  is injective. To see this, pick two different positive rationals  $x = p/q$  and  $y = r/s$  such that  $x \neq y$ . We need to prove  $g(x) \neq g(y)$ , that is,  $2^p 3^q \neq 2^r 3^s$ .

Since  $x \neq y$ , we have  $p \neq r$ , or  $q \neq s$ , or both. If  $p \neq r$ , then the largest power of 2 dividing  $g(x)$  and  $g(y)$  are different, implying  $g(x) \neq g(y)$ . If  $q \neq s$ , then the largest power of 3 dividing  $g(x)$  and  $g(y)$  are different, implying  $g(x) \neq g(y)$ . In either case,  $g(x) \neq g(y)$ . 

**Exercise:** Extend the above proof to give an injection  $g : \mathbb{Q} \rightarrow \mathbb{N}$ . Hint: use the fact that the union of two countable sets is countable. 

**Exercise:** What ordering of the (positive) rationals does the above give using the algorithm for getting ordering from the injective function? Order the first 7 positive rationals.

7. **The Set of Python Programs is Countable.**

Indeed, we show the set of strings  $\Sigma^*$  over any finite alphabet  $\Sigma$  is countable. Since  $\mathcal{P} \subseteq \Sigma^*$  for  $\Sigma$  given by all the  $< 200$  symbols on your keyboard, Theorem 1 would show  $\mathcal{P}$  is countable.

To do this, for any  $n \in \mathbb{N} \cup \{0\}$ , let us define  $\Sigma_n \subseteq \Sigma^*$  be the collection of all strings over the alphabet  $\Sigma$  which have *exactly* length  $n$ . Clearly,

$$\Sigma^* = \Sigma_0 \cup \Sigma_1 \cup \Sigma_2 \cup \dots = \bigcup_{n=0}^{\infty} \Sigma_n$$

*Observation.* For any fixed  $n$ , the set  $\Sigma_n$  is indeed a *finite* set. Indeed, it has size exactly  $|\Sigma|^n$  which is a large but finite number. And therefore, since finite sets are countable, there is at least one injective function

$$f_n : \Sigma_n \rightarrow \mathbb{N}$$

For instance, one could look at the *alphabetical ordering* of strings in  $\Sigma_n$ . This is well defined since  $\Sigma_n$  is finite.

And now we are ready to define the injective mapping  $h$  from  $\Sigma^*$  to  $\mathbb{N}$  using the same idea as in rationals. Given any  $\sigma \in \Sigma^*$ , define

$$h : \sigma \mapsto 2^{|\sigma|} \cdot 3^{f_n(\sigma)}$$

That is, if  $|\sigma| = n$  where  $n \in \mathbb{N} \cup \{0\}$ , then we map  $\sigma$  to  $2^n \cdot 3^{f_n(\sigma)}$ .

To see which this is an injection, let us select  $\sigma \neq \sigma'$  in  $\Sigma^*$ .

We claim this is an injection. To see this, take  $\sigma \neq \sigma'$ .

Case 1:  $|\sigma| \neq |\sigma'|$ . In this case the largest power of 2 dividing  $g(\sigma)$  and  $g(\sigma')$  are different, and thus the two numbers must be different.

Case 2:  $|\sigma| = |\sigma'| = n$ . In this case, both lie in  $\Sigma_n$  implying  $f_n(\sigma) \neq f_n(\sigma')$ . Thus, the largest power of 3 dividing  $g(\sigma)$  and  $g(\sigma')$  are different, and thus the two numbers must be different.

8. **Where are we headed?** The fact that  $\mathcal{P}$  is countable will lead us to the notion of “uncomputable” functions. What does that mean? For this we need to define what a computable function is. We will do so rather informally (and please take CS39 to get the rigorous version of computability) by saying

*A function  $f : \mathbb{N} \rightarrow \{0, 1\}$  is computable if there is a python code  $C$  taking input an number and outputting 0 or 1, such that for every  $n \in \mathbb{N}$ , we have  $C(n) = f(n)$ .*

**Theorem 4.** If every function in  $\mathcal{F}$  were computable, then  $\mathcal{F}$  would be a countable set.

*Proof.* We describe an injective map from  $\mathcal{F}$  to  $\mathcal{P}$ ; we would be done by Theorem 3.

Indeed, given a function  $f \in \mathcal{F}$ , since it is computable, there is a code  $C \in \mathcal{P}$  which computes it. We claim for two  $f \neq f' \in \mathcal{F}$  we can't have the same code  $C$ . Indeed, if  $f \neq f'$ , there exists some  $n \in \mathbb{N}$  such that  $f(n) \neq f'(n)$ . But both are  $C(n)$ . Contradiction.  $\square$

Next lecture, we show  $\mathcal{F}$  is *uncountable*. And thus, there must exist uncomputable functions.