

# Dissociation of Cognitive Effort–Based Decision Making and Its Associations With Symptoms, Cognition, and Everyday Life Function Across Schizophrenia, Bipolar Disorder, and Depression

Deanna M. Barch, Adam J. Culbreth, Dror Ben Zeev, Andrew Campbell, Subigya Nepal, and Erin K. Moran

## ABSTRACT

**BACKGROUND:** Anhedonia and amotivation are symptoms of many different mental health disorders that are frequently associated with functional disability, but it is not clear whether the same processes contribute to motivational impairments across disorders. This study focused on one possible factor, the willingness to exert cognitive effort, referred to as cognitive effort–cost decision making.

**METHODS:** We examined performance on the deck choice task as a measure of cognitive effort–cost decision making, in which people choose to complete an easy task for a small monetary reward or a harder task for larger rewards, in 5 groups: healthy control ( $n = 80$ ), schizophrenia/schizoaffective disorder ( $n = 50$ ), bipolar disorder with psychosis ( $n = 58$ ), current major depression ( $n = 60$ ), and past major depression ( $n = 51$ ). We examined cognitive effort–cost decision making in relation to clinician and self-reported motivation symptoms, working memory and cognitive control performance, and life function measured by ecological momentary assessment and passive sensing.

**RESULTS:** We found a significant diagnostic group  $\times$  reward interaction ( $F_{8,588} = 4.37, p < .001, \eta_p^2 = 0.056$ ). Compared with the healthy control group, the schizophrenia/schizoaffective and bipolar disorder groups, but not the current or past major depressive disorder groups, showed a reduced willingness to exert effort at the higher reward values. In the schizophrenia/schizoaffective and bipolar disorder groups, but not the major depressive disorder groups, reduced willingness to exert cognitive effort for higher rewards was associated with greater clinician-rated motivation impairments, worse working memory and cognitive control performance, and less engagement in goal-directed activities measured by ecological momentary assessment.

**CONCLUSIONS:** These findings suggest that the mechanisms contributing to motivational impairments differ among individuals with psychosis spectrum disorders versus depression.

<https://doi.org/10.1016/j.biopsych.2023.04.007>

Anhedonia and amotivation are clinical symptoms of many different mental health disorders that are frequently associated with distress, impairment, and functional disability (1). As an example, individuals with schizophrenia often report difficulties with motivation and anticipated pleasure/reward in the domains of social, occupational, and educational function that are associated with life impairment. In addition, individuals with major depression often report anhedonia (a reduction in the ability to experience pleasure) and a lack of motivation, again often associated with functional impairment (2–4). Furthermore, individuals with bipolar disorder can also experience anhedonia and amotivation, though they can experience hyper-reward responsivity in the manic phase (2). One hypothesis in the literature is that there are mechanisms contributing to anhedonia/amotivation that cut across putative diagnostic boundaries (5–9), though some work has begun to

suggest that such mechanisms may not be fully transdiagnostic (5,10–13). The goal of the current work was to test the hypothesis that transdiagnostic mechanisms contribute to motivational impairments, with a specific focus on the role of willingness to expend cognitive effort, a component of reward valuation in the positive valence systems component of the Research Domain Criteria (14–16).

There have been a number of different frameworks used to try to understand the pathways that lead individuals to experience anhedonia or amotivation (17). These frameworks include examining the experience of pleasure or reward in the moment, the anticipation of reward or pleasure in the future, and/or the ability to learn about cues and actions in the environment that might lead to reward or pleasure. One additional approach that may help integrate these different components of motivation and parse the source of impairments is the

concept of willingness to exert effort, or effort-cost decision making (ECDM). ECDM refers to choices that a person makes about how much effort to expend as a function of factors such as the amount or type of reward that one would receive, the likelihood of receiving that reward should one expend the effort, or the amount of time it would take to obtain the reward. Such effort can be either physical (e.g., finger tapping or grip exertion) or cognitive (e.g., willingness to perform a more difficult vs. less difficult cognitive task). There is now robust evidence that individuals with schizophrenia or other psychosis spectrum conditions display a reduced willingness to expend effort to obtain reward under certain conditions (high reward and high probability of obtaining the reward) that would typically lead people to allocate effort, both cognitive (18–22) and physical (20,21,23–32).

Studies investigating ECDM in depression have shown some evidence of reduced willingness to expend physical effort with increasing reward (25,33–37), consistent with the possibility that ECDM deficits represent a transdiagnostic mechanism contributing to amotivation and anhedonia. However, as with schizophrenia, there are mixed results, with some studies not finding impairments, even among those with current depression (23,24,38–40). A few studies have also examined cognitive effort allocation. Both Ang *et al.* (41) and Westbrook *et al.* (42) found that individuals with current depression discounted reward as a function of greater cognitive effort more so than healthy control participants (HCs), though Westbrook *et al.* found that those with remitted depression did not. In contrast, Vinckier *et al.* did not find that individuals with depression were less willing to allocate cognitive effort as a function of reward (38). Studies of ECDM in bipolar disorder have focused exclusively on physical effort, with evidence for reduced allocation of effort as a function of reward in depressed bipolar individuals in one study (37), but not in two other studies (24,43), in bipolar individuals with unspecified phase of illness (23,25), and in bipolar individuals during the manic phase of illness (24).

Several studies have also examined deficits in physical ECDM in transdiagnostic samples that include individuals with schizophrenia/schizoaffective disorder, major depressive disorder (MDD), and bipolar disorder. All 4 of these studies found that both individuals with schizophrenia/schizoaffective disorder and those with bipolar disorder (though with some variation for depressed vs. manic phase) demonstrated reduced willingness to allocate greater physical effort as a function of greater reward amount or probability (23–25,44), and Yang *et al.* and Zou *et al.* also found this to be true of individuals with current major depression (23,25). However, both Yang *et al.* (24) and Moran *et al.* (45) found that individuals with current major depression did not differ from HCs in their willingness to exert physical effort as a function of reward, and Yang *et al.* also did not see differences in patients with bipolar depression in a depressed episode. Furthermore, none of these studies examined cognitive effort allocation. Given that difficulties with concentration are a cardinal symptom of depression, it is possible that deficits in cognitive effort allocation might be more robust among individuals with current major depression.

Even if deficits in cognitive ECDM are present transdiagnostically, it is still possible that the mechanisms

contributing to cognitive ECDM deficits differ across putative diagnostic boundaries. For example, among individuals with schizophrenia, degree of working memory impairment was associated with effort discounting for cognitive effort (19). However, associations with cognitive function have not been examined in depression and bipolar disorder. Interestingly, there seems to be shared neural substrates of cognitive and physical ECDM in valuation regions that include parts of the dorsal frontal cortex, insula, and ventral striatum (46,47), but these different aspects of ECDM may dissociate in terms of their associations with more task-specific effector brain systems, such as frontal-parietal networks for cognitive function and motor/sensory regions for physical effort (46,47). As such, it is possible that the known disruptions in frontal-parietal systems found in psychosis (48) may also contribute to differential correlates of cognitive ECDM in psychosis versus mood disorders.

Furthermore, in schizophrenia, motivation and pleasure symptoms (e.g., anhedonia and amotivation) are associated with reduced effort allocation as a function of reward (either cognitive or physical) (19,21,24,27,28,31,49), though not in every study (26,30,39,50). In depression, reductions in anticipated pleasure have been associated with reduced effort allocation as a function of reward (33). In the physical effort transdiagnostic studies described above, Yang *et al.* (24) reported relationships of self-reported amotivation and effort allocation in schizophrenia, but not in depression or bipolar disorder. Neither Zou *et al.* (25) or Yang *et al.* found relationships of effort allocation with clinically assessed or self-reported symptoms. However, Moran *et al.* found relationships between self-reported amotivation/anhedonia that cut across schizophrenia, bipolar disorder, and major depression (44). Thus, whether associations with symptoms are transdiagnostic is still unclear. Furthermore, most studies examine relationships to clinician or self-reported symptoms, but not to everyday life experience measures with tools such as ecological momentary assessment (EMA) and passive sensing. In schizophrenia, greater willingness to exert physical and cognitive effort for reward was associated with fewer motivation and pleasure symptoms assessed via EMA (19,49,51). As such, determining whether laboratory-based measures of cognitive effort relate to everyday life experiences as assessed via EMA in patient populations is critical to understanding whether or not ECDM deficits are important in understanding functional impairment transdiagnostically. In theory, one could use traditional assessment measures (semistructured interviews and questionnaires) to gauge real-world function. However, there is evidence that these measures may be confounded by deficits in memory and associated cognitive processes in people with schizophrenia (49). EMA is an important tool for assessing daily experiences of motivation and pleasure in the moment, which might provide a more accurate assessment.

### Current Study

The goal of the current study was to examine whether there are transdiagnostic deficits in cognitive ECDM, a component of effort allocation that has not yet been examined in this way.

Cognitive effort could be particularly relevant to understanding life function because it may relate to ability to engage in educational and occupational endeavors, and it is not clear whether the transdiagnostic patterns of impairment in cognitive effort parallel those seen in physical effort. In addition, we wished to examine whether the symptom correlates of cognitive effort parallel those seen in physical effort. In addition, we wished to examine whether the symptom correlates of cognitive effort parallel those seen in physical effort. In addition, we wished to examine whether the symptom correlates of cognitive effort parallel those seen in physical effort. In addition, we wished to examine whether the symptom correlates of cognitive effort parallel those seen in physical effort.


## METHODS AND MATERIALS

### Participants

Study participants included 50 people with schizophrenia or schizoaffective disorder (SZ), 58 with bipolar I disorder (BD) with a history of psychosis in any current phase of illness, 60 with MDD as defined by the DSM-5 in a current major depressive episode (C-MDD), and 51 with a past history of a major depressive episode (P-MDD). We included both current and past depression groups to examine trait versus state associations with major depression. Additionally, there were 80 HCs. All participants provided written informed consent to the protocol approved by the Washington University Institutional Review Board. One hundred sixty-three of these individuals completed the tasks in person (40 SZ group, 44 BD group, 49 C-MDD group, and 51 HC group), and 87 (including all of the P-MDD group) completed them remotely via Zoom during COVID-19 in response to the temporary cessation of in-person testing. None of the results below differed as a function of mode of assessment (no interactions with assessment mode) (see the [Supplement](#)). Thus, all the data presented below used the entire sample. See the [Supplement](#) for inclusion/exclusion and additional study details.

### Diagnostic and Symptom Assessment

Diagnostic status, including history of psychosis in the individuals with BD, was confirmed using the Structured Clinical Interview for DSM-5 conducted by master's- or Ph.D.-level clinicians. These clinicians also assessed symptoms using the Structured Clinical Interview for DSM-4-TR Axis I disorders (52), the 24-item Brief Psychiatric Rating Scale (53–55), the Young Mania Rating Scale (56), the Bipolar Depression Rating Scale (57), and the Clinical Assessment Interview for Negative Symptoms (58–60). To assess motivation and pleasure negative symptoms, participants completed the Motivation and Pleasure Self-Report Scale (61) with higher scores indicating more motivation and pleasure across the week (i.e., lower motivation and pleasure negative symptoms). Self-reported depression symptoms were assessed using the Center for Epidemiological Studies Depression Scale (62), and mania



**Figure 1.** Illustration of the deck choice task.

symptoms were assessed using the Altman Self-Rating Mania Scale (45).

**Deck Choice Task.** We used the same cognitive effort-based decision-making task—the deck choice effort task—used previously in studies by Reddy *et al.* (20) and Horan *et al.* (21). In this task (Figure 1), participants make choices from one of two decks of cards. One deck of cards is all the same color (easy deck), and the participant does the same mental task for each card—deciding whether the number on the card is 1) odd or even or 2) above or below 5. The other deck of cards (hard deck) alternates between two colors, one of which has the individual do one mental task (odd/even) and the other a different task (above or below 5). Thus, modulation of cognitive effort in this paradigm is achieved via the amount of mental task switching a participant must perform during each deck. Participants complete practice rounds until they learn which color is associated with which task and achieve 70% accuracy. Participants earn a \$0.10 reward for successful completion of an easy deck choice, but hard deck choices can be associated with a \$0.10, \$0.20, or \$0.40 reward. There are 12 choices between easy and hard decks for each of the values of the hard deck (36 total), and participants do the easy or hard task 10 times for each choice. Successful completion is defined as at least 70% accuracy. Outcome measures are the percentage of hard tasks chosen at each of the 3 different reward amounts (\$0.10, \$0.20, and \$0.40). This task has reasonable reliability [intraclass correlation coefficient = 0.63 for change from low to high reward (20)].

**Dot Probe Expectancy Task.** This task is a measure of the goal representation component of cognitive control (63). See the [Supplement](#) for details.

**Running Span Task.** This task is a measure of working memory (64). See the [Supplement](#) for details.

### EMA and Passive Sensing

For the 163 individuals who completed the tasks in person, we used the Crosscheck (40) platform to conduct 2 weeks of EMA, along with passive sensing measures of movement during the

day. Crosscheck runs in the background on Android phones and passively collects data. The 87 virtual participants completed the exact same EMA component with the same timing and questions using Qualtrics surveys. Participants using Crosscheck also wore a Garmin Vivosmart 4 fitness watch on their nondominant arm to track steps, heart rate, and sleep throughout the 2-week EMA assessment period. We focused on daily step count as a measure of effort allocation for physical activity in the current analyses, though this measure was not available in the virtual participants.

For EMA, participants were prompted pseudorandomly 4 times a day between 9:30 AM and 8:30 PM and asked to answer questions about what they were currently doing and their current enjoyment and interest level and the same set of questions was asked about anticipated activities in the next 2 hours. We focused on the degree to which participants were currently engaged in putatively goal-directed activities as a measure potentially related to cognitive effort exertion. Goal directed was defined as choosing one or more of the following activities that required them to actively engage in an activity that accomplished some sort of occupational, work, or social task: 1) entertainment away from home, 2) exercising, 3) work/school, 4) running an errand, 5) cleaning/hygiene/chores/cooking, or 6) therapy/doctor's appointment. Nongoal directed was defined as choosing one or more of the following activities that could be passive: 1) eating or drinking, 2) television/radio/reading/computer, 3) socializing, 4) smoking, 5) sleeping, or 6) nothing in particular. Only EMA survey responses that were completed within 20 minutes of survey notification were included in the analyses. Mean response rates were 84% (SD = 18%) and did not differ by group (81%–86.5%;  $F_{4,295} = 0.833$ ,  $p = .51$ ). Consistent with previous EMA research (65), participants were only included in the analyses if they completed at least 33% of the surveys.

### Data Analysis

The deck choice task data were analyzed using a repeated-measures analysis of variance in SPSS version 27 that included diagnostic group (HC, SZ, BD, C-MDD, or P-MDD) as a between-subject factor and reward level (\$0.10, \$0.20, or \$0.40) as a within-subject factor. Significant diagnostic group  $\times$  reward level interactions were followed up by planned comparisons at each reward level. We computed correlations between the percentage of hard task choices in the high reward level (\$0.40) and performance on the working memory and cognitive control tasks, as well as the clinician-reported and self-report symptoms. Diagnostic group differences in correlations were compared using a Fisher  $r$ -to- $z$  transformation. The EMA and movement data were analyzed using linear mixed models in R version 4.0.3 using the lmer function in the package lme4 version 1.1.25. Because engagement in goal-directed activity was a binary outcome (1 or -1), we used a binary logistic linear mixed model for this analysis, using the glmer function in lme4. For these models, we used the percent choice of hard task in the \$0.40 reward condition and dummy codes for diagnostic group to predict either being engaged in a goal-directed activity when prompted, or steps taken, with the repeated estimates of the outcome variables across days nested as random effects within participants.

## RESULTS

### Demographic and Symptom Characteristics

There were no significant group differences in sex or age (Table 1). As expected, there were significant differences in education, with the SZ group having the lowest education; the HC, BD, and C-MDD groups having similar levels of education; and the P-MDD group having the highest level of education. There were also group differences in parental education, with HC having the lowest parental education, significantly lower than BD, C-MDD, and P-MDD, but not different from that of SZ.

### Group Differences

As shown in Figure 2, we observed a significant main effect of reward magnitude ( $F_{2,588} = 219.86$ ,  $p < .001$ ,  $\eta_p^2 = 0.48$ ), suggesting that people were more willing to expend cognitive effort as reward value increased. There was also a significant main effect of group ( $F_{4,294} = 3.82$ ,  $p = .005$ ,  $\eta_p^2 = 0.049$ ) that was modified by a significant group  $\times$  reward interaction ( $F_{8,588} = 4.37$ ,  $p < .001$ ,  $\eta_p^2 = 0.056$ ). Follow-up analyses revealed significant between group differences in hard cognitive task choice in the \$0.40 ( $F_{4,294} = 6.88$ ,  $p < .001$ ,  $\eta_p^2 = 0.086$ ) and \$0.20 reward conditions ( $F_{4,294} = 5.37$ ,  $p < .001$ ,  $\eta_p^2 = 0.068$ ), but no group differences in the \$0.10 reward condition ( $F_{4,294} = 1.05$ ,  $p = .383$ ,  $\eta_p^2 = 0.014$ ). For both the \$0.20 and \$0.40 reward conditions, these group differences reflected the SZ and BD groups making significantly fewer cognitive hard task choices than the HC, P-MDD, and C-MDD groups. The C-MDD and P-MDD groups did not make fewer cognitive hard task choices than HC in any condition.

### Relationships to Cognitive Task Difficulty

One reason that individuals with SZ or BD might be less likely to choose the hard cognitive task is that they could be less capable of performing that task. There were significant group differences in hard cognitive task success ( $F_{4,284} = 11.34$ ,  $p < .001$ ,  $\eta_p^2 = 0.138$ ). However, while individuals with SZ (mean = 48%, SD = 3.0%) were less able to complete the hard cognitive task than HCs (mean = 83%, SD = 2.4%) and those with BD (mean = 75%, SD = 3.3%), C-MDD (mean = 77%, SD = 3.0%), and P-MDD (mean = 79%, SD = 2.8%), the individuals with BD did not differ from those in the HC, C-MDD, or P-MDD groups in their ability to complete the hard cognitive task. Furthermore, when we included hard accuracy as a covariate in the analysis of variance, we still saw a significant group  $\times$  reward interaction ( $F_{8,566} = 2.15$ ,  $p < .05$ ,  $\eta_p^2 = 0.029$ ). We also found that the significant group  $\times$  reward interaction remained for both education ( $F_{8,586} = 2.32$ ,  $p < .05$ ,  $\eta_p^2 = 0.031$ ) and reading level ( $F_{8,584} = 2.99$ ,  $p = .003$ ,  $\eta_p^2 = 0.039$ ) as covariates. Thus, the deficits in willingness to exert effort for reward among the individuals with SZ and BD were not secondary to difficulties in performing the task or lower education or reading levels.

### Relationship to Working Memory and Cognitive Control

Because the SZ and BD groups performed similarly on the deck choice task, as did the C- and P-MDD groups, we combined each of these to create 3 groups (SZ/BD, C-/P-

**Table 1. Participant Demographics and Clinical Measures**

|   | HC, <i>n</i> = 80 | SZ, <i>n</i> = 50 | BD, <i>n</i> = 58 | C-MDD, <i>n</i> = 60 | P-MDD, <i>n</i> = 51 | Group Differences, <i>p</i> Value | Pattern                               |
|---|-------------------|-------------------|-------------------|----------------------|----------------------|-----------------------------------|---------------------------------------|
| Sex, Female:Male                        | 47:33             | 28:22             | 37:21             | 37:23                | 29:22                | .75                               | –                                     |
| <b>Race</b>                             |                   |                   |                   |                      |                      |                                   |                                       |
| African American                        | 39                | 31                | 10                | 22                   | 10                   | <.001                             | –                                     |
| Caucasian                               | 37                | 19                | 45                | 37                   | 39                   | <.001                             | –                                     |
| Additional races                        | 4                 | 0                 | 3                 | 1                    | 2                    | <.001                             | –                                     |
| Age, Years                              | 36.39 (9.19)      | 35.83 (8.51)      | 37.56 (8.78)      | 33.63 (8.71)         | 36.45 (8.44)         | .16                               | –                                     |
| Education, Years                        | 15.58 (2.38)      | 13.58 (2.30)      | 15.09 (2.51)      | 15.07 (2.24)         | 16.49 (2.43)         | <.001                             | P-MDD > HC = BD = C-MDD > SZ          |
| Parental Education, Years               | 13.58 (2.75)      | 14.28 (3.02)      | 14.79 (3.09)      | 15.17 (2.69)         | 16.06 (2.98)         | <.001                             | P-MDD > HC = SZ = BD; BD = C-MDD > HC |
| WTAR                                    | 37.04 (9.26)      | 31.04 (12.06)     | 39.67 (8.76)      | 37.17 (9.56)         | 41.72 (7.42)         | <.001                             | P-MDD > HC, BD, C-MDD > SZ            |
| Unmedicated                             | 100%              | 20%               | 16%               | 29%                  | 50%                  | –                                 | –                                     |
| Antidepressant                          | 0                 | 52%               | 71%               | 68%                  | 48%                  | –                                 | –                                     |
| Mood Stabilizer                         | 0                 | 14%               | 59%               | 7%                   | 6%                   | –                                 | –                                     |
| Antipsychotic                           | 0                 | 80%               | 36%               | 10%                  | 4%                   | –                                 | –                                     |
| CPZ                                     | –                 | 481.2 (263.0)     | 342.7 (212.5)     | 167.7 (111.7)        | 175.0 (35.4)         | –                                 | –                                     |
| Dot Probe Expectancy <i>d'</i> -Context | 2.62 (1.25)       | 1.58 (1.09)       | 2.27 (1.16)       | 2.52 (0.97)          | 2.37 (1.26)          | <.001                             | HC = BD = C-MDD = P-MDD > SZ          |
| Running Span Total Correct              | 47.83 (15.4)      | 34.43 (15.9)      | 44.36 (17.2)      | 43.82 (14.8)         | –                    | .002                              | HC = BD = C-MDD > SZ                  |
| CAINS—Motivation and Pleasure Deficits  | –                 | 14.50 (7.74)      | 10.98 (6.35)      | 14.32 (5.59)         | 9.45 (5.42)          | <.001                             | SZ = C-MDD > BD = P-MDD               |
| BPRS Depression                         | –                 | 10.06 (4.03)      | 9.81 (4.39)       | 13.48 (3.20)         | 9.37 (3.57)          | <.001                             | C-MDD > SZ = BD = P-MDD               |
| BPRS Mania                              | –                 | 7.58 (2.79)       | 7.76 (4.30)       | 5.53 (0.91)          | 6.00 (1.15)          | <.001                             | SZ = BD > C-MDD = P-MDD               |
| BPRS Negative                           | –                 | 8.04 (2.79)       | 6.05 (2.38)       | 6.62 (2.27)          | 6.16 (2.17)          | <.001                             | SZ > BD = C-MDD = P-MDD               |
| BPRS Positive                           | –                 | 9.56 (4.31)       | 4.21 (2.25)       | 3.55 (1.10)          | 3.35 (0.96)          | <.001                             | SZ > BD = C-MDD = P-MDD               |
| CES-D                                   | 5.48 (3.46)       | 12.06 (6.66)      | 11.64 (6.34)      | 16.08 (5.04)         | 9.63 (5.33)          | <.001                             | C = MDD > SZ = BD > P-MDD > HC        |
| MAP-SR                                  | 41.95 (9.86)      | 33.57 (12.46)     | 36.28 (9.74)      | 28.85 (9.59)         | 36.80 (8.87)         | <.001                             | C-MDD < SZ < BD = P-MDD < HC          |
| ASRM                                    | 4.72 (4.18)       | 4.60 (3.12)       | 4.72 (4.13)       | 2.80 (2.87)          | 2.76 (2.96)          | .001                              | HC = SZ = BD > C-MDD = P-MDD          |


Values are provided as *n*, mean (SD), or %. Higher MAP-SR scores indicate more motivation and pleasure across the week (i.e., lower motivation and pleasure negative symptoms).

ASRM, Altman Self-Rating Mania Scale; BD, bipolar disorder; BPRS, Brief Psychiatric Rating Scale; CAINS, Clinical Assessment Interview for Negative Symptoms; CES-D, Center for Epidemiological Studies Depression Scale; C-MDD, current major depressive disorder; CPZ, chlorpromazine-equivalent dose; HC, healthy control participant; MAP-SR, Motivation and Pleasure Self-Report; MDD, major depressive disorder; P-MDD, past major depressive disorder; SZ, schizophrenia; WTAR, Wechsler Test of Adult Reading.

MDD, and HC) for the correlational analyses. As shown in Table 2, running span task performance as a measure of working memory was significantly positively associated with the percentage of hard task choices in the highest reward conditions in SZ/BD, but not in HC or C-/P-MDD groups, though the magnitude of these correlations did not differ significantly across groups. *d'*-context on the dot probe expectancy task as a measure of cognitive control was significantly positively associated with the percentage of hard task choices in the highest reward conditions in both the HC and SZ/BD groups, but not in the MDD group. Furthermore, the magnitude of the association was significantly stronger in the SZ/BD group compared with the MDD group (Table 2).

**Relationships With Symptoms**

As shown in Table 2, among the SZ/BD group, clinician-rated motivation and pleasure symptoms were negatively associated



**Figure 2.** Percentage of hard cognitive task choices across groups. Error bars reflect 95% CIs. BD, bipolar disorder; C-MDD, current major depressive disorder; HC, healthy control participant; P-MDD, past major depressive disorder; SZ, schizophrenia.

**Table 2. Correlations With Percentage or Hard Task Choices in the Highest Reward Condition (\$0.40)**

| Measure   | Correlations      |                    |                 | Fisher's <i>r</i> -to- <i>z</i> Transformations Comparing Correlations Across Groups |                    |                        |
|---|-------------------|--------------------|-----------------|--|--------------------|------------------------|
|   | HC                | SZ and BD          | C-MDD and P-MDD | HC vs. SZ/BD   | HC vs. C-MDD/P-MDD | SZ/BD vs. C-MDD/P-MDD  |
| Running Span Percent Correct                        | 0.22              | 0.26 <sup>a</sup>  | 0.15            | Z = 0.22   | Z = 0.34           | Z = 0.60               |
| Dot Probe Expectancy <i>d'</i> -Context             | 0.26 <sup>a</sup> | 0.37 <sup>b</sup>  | 0.18            | Z = 0.77   | Z = 1.24           | Z = 2.19 <sup>a</sup>  |
| CAINS Motivation and Pleasure                       | NA                | -0.29 <sup>a</sup> | -0.06           | NA   | NA                 | Z = -1.74 <sup>a</sup> |
| BPRS Depression                                     | NA                | -0.09              | 0.08            | NA   | NA                 | Z = -1.24              |
| Motivation and Pleasure Self-Report                 | 0.14              | 0.10               | 0.001           | Z = -0.33  | Z = 1.00           | Z = 0.72               |
| Center for Epidemiological Studies Depression Scale | 0.03              | 0.05               | 0.07            | Z = 0.13   | Z = -0.26          | Z = -0.15              |

Higher scores on the Motivation and Pleasure Self-Report Scale indicate more motivation and pleasure across the week (i.e., lower motivation and pleasure negative symptoms).

BD, bipolar disorder; BPRS, Brief Psychiatric Rating Scale; CAINS, Clinical Assessment Interview for Negative Symptoms; C-MDD, current major depressive disorder; HC, healthy control participant; NA, not applicable; P-MDD, past major depressive disorder; SZ, schizophrenia.

<sup>a</sup>*p* < .05.

<sup>b</sup>*p* < .01.

with the percentage of hard task choices in SZ/BD, but not in C-/P-MDD. This relationship was significantly stronger in the SZ/BD compared with the MDD group. Clinician-rated depression symptoms were not related to the percentage of hard task choices in either group. Self-reported motivation and pleasure symptoms and depression were also not related to cognitive hard task choice in any group (Table 2).

### Relationships to Engagement in Goal-Directed Activities or Locomotion

The binary logistic linear mixed effect model predicting the likelihood of being engaged in a goal-directed activity when questioned via EMA indicated a significant positive relationship to the percent of hard task choices at the \$0.40 reward level (standardized coefficient [SDCoef.] = 0.19, *z* = 3.54, *p* = .0004). When we added the 3 group factor (HC, SZ/BD, C-/P-MDD) to the glmer model, the relationship of percent of hard task choices at the \$0.40 reward level to likelihood of being engaged in goal-directed activities remained significant (SDCoef. = 0.11, *z* = 2.11, *p* = .035). Interestingly, both patient groups were less likely to be engaged in goal-directed activities than HCs, though the magnitude of this difference was larger for the SZ/BD (SDCoef. = -0.33, *z* = -5.27, *p* < .0001) than the MDD (SDCoef. = -0.14, *t*<sub>289</sub> = -2.29, *p* = .022) group. A model that included interactions between the group factor and percent hard task choices in predicting goal-directed activity fit the data worse (Bayesian information criterion = 17,876.3 for interaction model vs. 17,859.9 without), and neither interaction term was significant (*p* > .50). When we examined this relationship within each group, it was only significant in the SZ/BD group (SDCoef. = 0.19, *z* = 2.618, *p* < .009) and not in the HC (SDCoef. = 0.06, *z* = .71, *p* = .479) or MDD groups (SDCoef. = -0.01, *z* = -0.135, *p* = .893). Thus, task performance and diagnostic group status demonstrated independent relationships with engagement in goal-directed activities, with some evidence of stronger relations in SZ/BD.

There was also a significant positive relationship between percent of hard task choices and number of steps taken during the day (SDCoef. = 0.22, *z* = 2.45, *p* = .0155). This effect remained significant (SDCoef. = 0.20, *z* = 2.10, *p* = .0369) in a

model that included the group factor, and neither group differed from the HC group in the number of steps taken per day. There were, again, no significant interactions between group and percent hard task choices in relationship to steps taken (*ps* > .80), and, in this case, none of the within-group associations was significant, though the closest was the SZ/BD group (SDCoef. = 0.22, *z* = 1.68, *p* = .0974).

### DISCUSSION

We found evidence of reduced willingness to exert cognitive effort compared with HCs among individuals with SZ and BD with psychosis, but not among those with C- or P-MDD. We also found that reduced cognitive ECDM was related to impaired working memory and cognitive control and to clinician-rated symptoms of reduced motivation and pleasure in individuals with SZ and BD, but not among those with C- or P-MDD. Furthermore, willingness to exert cognitive effort in the laboratory-based task was related to the likelihood of being engaged in goal-directed activities in everyday life via EMA in the SZ/BD group, but not in the C-/P-MDD group, though the average number of steps taken during the day was related across all groups. There was also a significant reduction in goal-directed activities among all diagnostic groups compared with the HC group, with the largest differences among individuals with SZ and BD. Together, these findings challenge the hypothesis that the same mechanisms contribute to motivational impairments across SZ, BD, and MDD, instead suggesting commonalities across SZ and BD with psychosis that may not extend to MDD.

Our findings demonstrate that both individuals with SZ and those with BD with psychosis show a reduced willingness to exert cognitive effort as a function of reward compared with HCs, with the largest group differences in the highest reward condition. This pattern in SZ replicates many previous findings with both cognitive (18–22) and physical (20,21,23–32) effort allocation. However, no previous research has examined cognitive effort allocation in a BD group. Our findings in BD are consistent with several previous studies that found reduced willingness to exert physical effort as a function of reward (23–25,37) but contrast with 2 other studies that did not find

such a reduction (24,43). One intriguing potential cause of the variability across studies is whether the individuals with BD had psychosis. All the individuals with BD in the current study had a history of psychosis. Hershenberg *et al.* (37), Wang *et al.* (23), and Zou *et al.* (25) did not specify whether the individuals with BD in their studies had psychosis, and they all observed deficits in willingness to exert physical effort. Yang *et al.* (24) found deficits in individuals with BD in a manic phase, 10 of whom had psychosis. However, all individuals with BD in the Whitton *et al.* (43) study had psychosis but did not show physical effort allocation deficits. Notably, however, Whitton *et al.* never presented demographic or clinical information separately as a function of diagnostic group, so it is difficult to know whether there may have been confounding factors that might have affected the findings. Nonetheless, the fact that the individuals with SZ and BD with psychosis showed very similar patterns of cognitive ECDM deficits is consistent with a large body of literature, suggesting commonalities across these putatively different disorders in terms of genetics, neurobiology, and cognitive function (64,66–71) and providing evidence consistent with the idea that BD with psychosis may be part of a spectrum of psychotic disorders.

In striking contrast to the findings in SZ and BD with psychosis, neither the individuals with current depression nor those with past depression showed any evidence of a reduced willingness to exert cognitive effort. It is difficult to say whether this is consistent with the prior literature on either physical or cognitive effort allocation in depression because the previous work in this area has been mixed, with several studies finding physical ECDM deficits in depression (25,33–37) but others not finding such deficits (23,24,38,39). Work on cognitive ECDM has been similarly mixed, with 2 studies finding greater discounting of reward as a function of cognitive effort in depression (41,42), but other work (38) finding no differences in willingness to exert cognitive effort as a function of reward. As with the mixed findings in BD, it is not clear what drives the variability in findings across studies of depression. One possibility is that it could have something to do with an aspect of depressive disorder severity that has not yet been systematically examined, such as age of onset, number of prior episodes, or comorbid anxiety, hypotheses that could be tested in future studies.

In the current study, the individuals with current depression showed similar levels of clinician-rated symptoms of reduced motivation and pleasure to those of individuals with SZ and more symptoms than individuals with BD. Furthermore, they self-reported worse symptoms of motivation and pleasure than individuals with either SZ or BD (who did not differ). Thus, the differences at the group level between individuals with SZ/BD and MDD in the current study cannot easily be explained by less severe symptoms of impaired motivation in C-MDD compared with the other groups. Furthermore, when we examined the relationship of clinician-rated motivation and pleasure symptoms to ECDM, there was a relationship in the SZ/BD group, but not the C-/P-MDD group, and this relationship was significantly greater in SZ/BD than in MDD. We also found that cognitive ECDM performance was related to working memory and cognitive control performance in the SZ/BD, but not in the MDD group, again with significantly stronger associations in the SZ/BD group (at least for cognitive control). Together these results are consistent with the idea that the

mechanisms contributing to deficits in motivation differ across individuals in the psychosis spectrum versus MDD. Furthermore, they suggest that ECDM deficits play a more important role in motivational impairments in the psychosis spectrum than in depression, with potential contributions from cognitive impairment. The latter hypothesis is consistent with the fact that there tends to be associations between negative symptoms in psychosis (which include motivation impairments) and cognitive dysfunction (72).

The data hint that the relationship of cognitive ECDM to everyday life function as measured by EMA differed between the psychosis spectrum and depression groups. We found that the number of hard task choices in the highest reward value predicted greater likelihood of being engaged in goal-directed activities at the time of EMA prompts. While the diagnostic group also predicted reduced engagement in goal-directed activities, particularly among the SZ/BD group, the relation of cognitive effort exertion remained independent of group status. There was no significant modulation of this relationship by group, but the association of cognitive ECDM performance and engagement in goal-directed activities was only significant in the SZ/BD group and nonexistent in the MDD group. We also found that better ECDM task performance predicted more daily steps taken on average. This association was again independent of group status, with no group differences in steps taken, though, again, data hinted at a stronger relationship in the SZ/BD group. Taken together, these findings provide support for a link between cognitive ECDM and function in everyday life, for both engagement in goal-directed activities and steps taken, which are intriguingly likely to involve both cognitive and physical effort exertion. These findings also provide some evidence consistent with this relationship's being different in the SZ/BD than the MDD group, at least for goal-directed activities. In future work, it will be important to provide even more detailed examination of the nature of goal-directed activities (what kinds of work, what type of schooling, etc.) being performed in everyday life, as this may reveal even stronger evidence for diagnostic group differences in the relationship between willingness to exert cognitive effort and function in everyday life.

These findings must be interpreted in the context of several limitations. The data are cross-sectional, and longitudinal analyses of the variation in laboratory-based assessments and function in everyday life would help to establish the robustness of such relationships. In addition, we did not have a BD group without a history of psychosis or an MDD group with psychosis. Thus, we cannot fully assess the degree to which the presence of psychosis is critical in determining the presence of relationships between cognitive ECDM and symptoms of motivational impairment and measures of function in everyday life. Furthermore, a subset of the participants completed the tasks virtually (including all the individuals with P-MDD), though we did not find that the results differed meaningfully as a function of in-person versus virtual assessment, and the individuals with C- and P-MDD performed almost identically. Finally, we found relationships to clinician-rated motivation and pleasure symptoms, but not to the same domain of self-report symptoms. Further work will be needed to determine why results differ across clinician versus self-report, and whether one or the other might be more predictive of different domains of motivation-relevant behaviors.

In summary, the current study provides strong evidence for the differential presence of deficits in cognitive ECDM among individuals with psychosis spectrum disorders (but not among individuals with current or past depression). Furthermore, we found evidence for differential relationships of cognitive ECDM to symptoms, cognitive function, and everyday life function as measured by EMA among individuals with psychosis spectrum disorders, but not among those with depression. These differential relationships suggest that the mechanisms contributing to motivational impairments are not fully transdiagnostic and differ across the psychosis versus nonpsychosis divide.

### ACKNOWLEDGMENTS AND DISCLOSURES

This work was funded by National Institute of Mental Health (Grant Nos. R01MH066031 [to DMB, EKM, SN, DBZ, AC] and K23126986 [to AJC]).

We thank the participants in this study who gave generously of their time. We also thank those who helped with all aspects of data collection.

Aside from funding for this project, DMB receives funding from the National Institute on Drug Abuse and the American Foundation for Suicide Prevention and consults for Cerevence. DBZ has financial interests in Merlin LLC, FOCUS technology, and CORE technology; has an intervention content licensing agreement with Pear Therapeutics; and has provided consultation services to Trusst Health, K Health, Boehringer Ingelheim, eQuility, Deep Valley Labs, and Otsuka Pharmaceuticals. All other authors report no biomedical financial interests or potential conflicts of interest.

### ARTICLE INFORMATION

From the Department of Psychological & Brain Sciences, Washington University in St. Louis, St. Louis, Missouri (DMB, EKM); Department of Psychiatry, Washington University School of Medicine, St. Louis, Missouri (DMB); Department of Radiology, Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, Missouri (DMB); Maryland Psychiatric Research Center, Department of Psychiatry, University of Maryland, Baltimore, Maryland (AJC); Department of Psychiatry, University of Washington, Seattle, Washington (DBZ); and Department of Computer Science, Dartmouth College, Hanover, New Hampshire (AC, SN).

Address correspondence to Deanna M. Barch, Ph.D., at [dbarch@wustl.edu](mailto:dbarch@wustl.edu).

Received Jan 25, 2023; revised Apr 5, 2023; accepted Apr 6, 2023.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.biopsych.2023.04.007>.

### REFERENCES

- Trøstheim M, Eikemo M, Meir R, Hansen I, Paul E, Kroll SL, *et al.* (2020): Assessment of anhedonia in adults with and without mental illness: A systematic review and meta-analysis. *JAMA Netw Open* 3: e2013233.
- Nusslock R, Alloy LB (2017): Reward processing and mood-related symptoms: An RDoC and translational neuroscience perspective. *J Affect Disord* 216:3–16.
- Craske MG, Meuret AE, Ritz T, Treanor M, Dour HJ (2016): Treatment for anhedonia: A neuroscience driven approach. *Depress Anxiety* 33:927–938.
- Fletcher K, Parker G, Paterson A, Fava M, Iosifescu D, Pizzagalli DA (2015): Anhedonia in melancholic and non-melancholic depressive disorders. *J Affect Disord* 184:81–88.
- Whitton AE, Treadway MT, Pizzagalli DA (2015): Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Curr Opin Psychiatry* 28:7–12.
- Zhang B, Lin P, Shi H, Öngür D, Auerbach RP, Wang X, *et al.* (2016): Mapping anhedonia-specific dysfunction in a transdiagnostic approach: An ALE meta-analysis. *Brain Imaging Behav* 10:920–939.
- Guineau MG, Ikani N, Rinck M, Collard RM, van Eijndhoven P, Tendolkar I, *et al.* (2022): Anhedonia as a transdiagnostic symptom across psychological disorders: A network approach. *Psychol Med* 53(9):1–12.
- Gooding DC, Pflum M (2022): The transdiagnostic nature of social anhedonia: Historical and current perspectives. *Curr Top Behav Neurosci* 58:381–395.
- Schaub AC, Kirschner M, Schweinfurth N, Mähmann L, Kettelhack C, Engeli EE, *et al.* (2021): Neural mapping of anhedonia across psychiatric diagnoses: A transdiagnostic neuroimaging analysis. *NeuroImage Clin* 32: 102825.
- Lambert C, Da Silva S, Geniti AK, Rizvi SJ, Foussias G, Kennedy SH (2018): Anhedonia in depression and schizophrenia: A transdiagnostic challenge. *CNS Neurosci Ther* 24:615–623.
- Strauss GP, Cohen AS (2017): A transdiagnostic review of negative symptom phenomenology and etiology. *Schizophr Bull* 43:712–719.
- Culbreth AJ, Moran EK, Barch DM (2018): Effort-cost decision-making in psychosis and depression: Could a similar behavioral deficit arise from disparate psychological and neural mechanisms? *Psychol Med* 48:889–904.
- Barch DM, Pagliaccio D, Luking K (2016): Mechanisms underlying motivational deficits in psychopathology: Similarities and differences in depression and schizophrenia. *Curr Top Behav Neurosci* 27:411–449.
- Morris SE, Cuthbert BN (2012): Research Domain Criteria: Cognitive systems, neural circuits, and dimensions of behavior. *Dialogues Clin Neurosci* 14:29–37.
- Der-Avakian A, Pizzagalli DA (2018): Translational assessments of reward and anhedonia: A tribute to Athina Markou. *Biol Psychiatry* 83:932–939.
- Hess JL, Kawaguchi DM, Wagner KE, Faraone SV, Glatt SJ (2016): The influence of genes on “positive valence systems” constructs: A systematic review. *Am J Med Genet B Neuropsychiatr Genet* 171B:92–110.
- Barch DM, Gold JM, Kring AM (2017): Paradigms for assessing hedonic processing and motivation in humans: Relevance to understanding negative symptoms in psychopathology. *Schizophr Bull* 43:701–705.
- Chang WC, Westbrook A, Strauss GP, Chu AOK, Chong CSY, Siu CMW, *et al.* (2020): Abnormal cognitive effort allocation and its association with amotivation in first-episode psychosis. *Psychol Med* 50:2599–2609.
- Culbreth AJ, Moran EK, Kandala S, Westbrook A, Barch DM (2020): Effort, avolition and motivational experience in schizophrenia: Analysis of behavioral and neuroimaging data with relationships to daily motivational experience. *Clin Psychol Sci* 8:555–568.
- Reddy LF, Horan WP, Barch DM, Buchanan RW, Dunayevich E, Gold JM, *et al.* (2015): Effort-based decision-making paradigms for clinical trials in schizophrenia: Part 1—psychometric characteristics of 5 paradigms. *Schizophr Bull* 41:1045–1054.
- Horan WP, Reddy LF, Barch DM, Buchanan RW, Dunayevich E, Gold JM, *et al.* (2015): Effort-based decision-making paradigms for clinical trials in schizophrenia: Part 2—external validity and correlates. *Schizophr Bull* 41:1055–1065.
- Gold JM, Kool W, Botvinick MM, Hubzin L, August S, Waltz JA (2015): Cognitive effort avoidance and detection in people with schizophrenia. *Cogn Affect Behav Neurosci* 15:145–154.
- Wang YY, Wang Y, Huang J, Sun XH, Wang XZ, Zhang SX, *et al.* (2022): Shared and distinct reward neural mechanisms among patients with schizophrenia, major depressive disorder, and bipolar disorder: An effort-based functional imaging study. *Eur Arch Psychiatry Clin Neurosci* 272:859–871.
- Yang X, Huang J, Harrison P, Roser ME, Tian K, Wang D, Liu G (2021): Motivational differences in unipolar and bipolar depression, manic bipolar, acute and stable phase schizophrenia. *J Affect Disord* 283:254–261.
- Zou YM, Ni K, Wang YY, Yu EQ, Lui SSS, Zhou FC, *et al.* (2020): Effort-cost computation in a transdiagnostic psychiatric sample: Differences among patients with schizophrenia, bipolar disorder, and major depressive disorder. *Psych J* 9:210–222.
- Huang J, Yang XH, Lan Y, Zhu CY, Liu XQ, Wang YF, *et al.* (2016): Neural substrates of the impaired effort expenditure decision making in schizophrenia. *Neuropsychology* 30:685–696.



27. Strauss GP, Bartolomeo LA, Luther L (2023): Reduced willingness to expend effort for rewards is associated with risk for conversion and negative symptom severity in youth at clinical high-risk for psychosis. *Psychol Med* 53:714–721.
28. Cooper JA, Barch DM, Reddy LF, Horan WP, Green MF, Treadway MT (2019): Effortful goal-directed behavior in schizophrenia Computational subtypes and associations with cognition. *J Abnorm Psychol* 128:710–722.
29. Barch DM, Treadway MT, Schoen N (2014): Effort, anhedonia, and function in schizophrenia: Reduced effort allocation predicts amotivation and functional impairment. *J Abnorm Psychol* 123:387–397.
30. Fervaha G, Graff-Guerrero A, Zakzanis KK, Foussias G, Agid O, Remington G (2013): Incentive motivation deficits in schizophrenia reflect effort computation impairments during cost–benefit decision-making. *J Psychiatr Res* 47:1590–1596.
31. Gold JM, Strauss GP, Waltz JA, Robinson BM, Brown JK, Frank MJ (2013): Negative symptoms of schizophrenia are associated with abnormal effort-cost computations. *Biol Psychiatry* 74:130–136.
32. Chang WC, Chu AOK, Treadway MT, Strauss GP, Chan SKW, Lee EHM, *et al.* (2019): Effort-based decision-making impairment in patients with clinically stabilized first-episode psychosis and its relationship with amotivation and psychosocial functioning. *Eur Neuropsychopharmacol* 29:629–642.
33. Yang XH, Huang J, Zhu CY, Wang YF, Cheung EF, Chan RC, Xie GR (2014): Motivational deficits in effort-based decision making in individuals with subsyndromal depression, first-episode and remitted depression patients. *Psychiatry Res* 220:874–882.
34. Berwian IM, Wenzel JG, Collins AGE, Seifritz E, Stephan KE, Walter H, Huys QJM (2020): Computational mechanisms of effort and reward decisions in patients with depression and their association with relapse after antidepressant discontinuation. *JAMA Psychiatry* 77:513–522.
35. Treadway MT, Bossaller NA, Shelton RC, Zald DH (2012): Effort-based decision-making in major depressive disorder: A translational model of motivational anhedonia. *J Abnorm Psychol* 121:553–558.
36. Treadway MT, Buckholz JW, Schwartzman AN, Lambert WE, Zald DH (2009): Worth the ‘‘Effort’’? The effort expenditure for rewards task as an objective measure of motivation and anhedonia. *PLoS One* 4: e6598.
37. Hershenberg R, Satterthwaite TD, Dalal A, Katchmar N, Moore TM, Kable JW, Wolf DH (2016): Diminished effort on a progressive ratio task in both unipolar and bipolar depression. *J Affect Disord* 196:97–100.
38. Vinckier F, Jaffre C, Gauthier C, Smajda S, Abdel-Ahad P, Le Bouc R, *et al.* (2022): Elevated effort cost identified by computational modeling as a distinctive feature explaining multiple behaviors in patients with depression. *Biol Psychiatry Cogn Neurosci Neuroimaging* 7:1158–1169.
39. Cathomas F, Klaus F, Guetter K, Seifritz E, Hartmann-Riemer MN, Tobler PN, *et al.* (2021): Associations between negative symptoms and effort discounting in patients with schizophrenia and major depressive disorder. *Schizophrenia Bulletin Open* 2:sgab022.
40. Wang R, Aung MSH, Abdullah S, Brian R, Campbell AT, Choudhury T, *et al.* (2016): CrossCheck: Toward passive sensing and detection of mental health changes in people with schizophrenia. In: *Proceedings of the 2016 ACM International Joint Conference on Pervasive and Ubiquitous Computing*. New York, NY: Association for Computing Machinery, 886–897.
41. Ang Y-S, Gelda SE, Pizzagalli DA (2022): Cognitive effort-based decision-making in major depressive disorder [published online April 25]. *Psychol Med*.
42. Westbrook A, Yang X, Bylsma LM, Daches S, George CJ, Seidman AJ, *et al.* (2022): Economic choice and heart rate fractal scaling indicate that cognitive effort is reduced by depression and boosted by sad mood. *Biol Psychiatry Cogn Neurosci Neuroimaging*.
43. Whitton AE, Merchant JT, Lewandowski KE (2020): Dissociable mechanisms underpinning effort-cost decision-making across the psychosis spectrum. *Schizophr Res* 224:133–140.
44. Moran EK, Prevost C, Culbreth AJ, Barch DM (2023): Effort-based decision-making in psychotic and mood disorders. *J Psychopathol Clin Sci* 132:490–498.
45. Altman EG, Hedeker D, Peterson JL, Davis JM (1997): The Altman self-rating mania scale. *Biol Psychiatry* 42:948–955.
46. Schmidt L, Lebreton M, Cléry-Melin ML, Daunizeau J, Pessiglione M (2012): Neural mechanisms underlying motivation of mental versus physical effort. *PLoS Biol* 10:e1001266.
47. Chong TT, Apps M, Giehl K, Silence A, Grima LL, Husain M (2017): Neurocomputational mechanisms underlying subjective valuation of effort costs. *PLoS Biol* 15:e1002598.
48. Minzenberg MJ, Laird AR, Thelen S, Carter CS, Glahn DC (2009): Meta-analysis of 41 functional neuroimaging studies of executive function in schizophrenia. *Arch Gen Psychiatry* 66:811–822.
49. Moran EK, Culbreth AJ, Barch DM (2017): Ecological momentary assessment of negative symptoms in schizophrenia: Relationships to effort-based decision making and reinforcement learning. *J Abnorm Psychol* 126:96–105.
50. Docx L, de la Asuncion J, Sabbe B, Hoste L, Baeten R, Warnaeys N, Morrens M (2015): Effort discounting and its association with negative symptoms in schizophrenia. *Cogn Neurosci* 20:172–185.
51. Culbreth AJ, Westbrook A, Braver TS, Barch DM (2020): Effort in daily life: Relationships between experimental tasks and daily experience. *Motiv Sci* 6:303–308.
52. First MB, Williams JBW, Karg RS, Spitzer RL (2016): User’s guide for the SCID-5-CV Structured Clinical Interview for DSM-5® disorders: Clinical version. Washington, DC: American Psychiatric Publishing, Inc.
53. Ventura J, Green MF, Shaner A, Liberman RP (1993): Training and quality assurance on the Brief Psychiatric Rating Scale: The ‘‘drift busters’’. *Int J Methods Psychiatry Res* 3:221–226.
54. Ventura J, Lukoff D, Nuechterlein KH, Liberman RP, Green MF, Shaner A (1993): Brief Psychiatric Rating Scale (BPRS) expanded version: Scales, anchor points, and administration manual. *Int J Psychiatr Methods* 3:227–243.
55. Overall JE, Gorham DR (1962): The brief psychiatric rating scale. *Psychol Rep* 10:799–812.
56. Young RC, Biggs JT, Ziegler VE, Meyer DA (1978): A rating scale for mania: Reliability, validity and sensitivity. *Br J Psychiatry* 133:429–435.
57. Berk M, Malhi GS, Cahill C, Carman AC, Hadzi-Pavlovic D, Hawkins MT, *et al.* (2007): The Bipolar Depression Rating Scale (BDRS): Its development, validation and utility. *Bipolar Disord* 9:571–579.
58. Forbes C, Blanchard JJ, Bennett M, Horan WP, Kring A, Gur R (2010): Initial development and preliminary validation of a new negative symptom measure: The Clinical Assessment Interview for Negative Symptoms (CAINS). *Schizophr Res* 124:36–42.
59. Blanchard JJ, Kring AM, Horan WP, Gur R (2011): Toward the next generation of negative symptom assessments: The collaboration to advance negative symptom assessment in schizophrenia. *Schizophr Bull* 37:291–299.
60. Horan WP, Kring AM, Gur RE, Reise SP, Blanchard JJ (2011): Development and psychometric validation of the Clinical Assessment Interview for Negative Symptoms (CAINS). *Schizophr Res* 132:140–145.
61. Llerena K, Park SG, McCarthy JM, Couture SM, Bennett ME, Blanchard JJ (2013): The Motivation and Pleasure Scale-Self-Report (MAP-SR): Reliability and validity of a self-report measure of negative symptoms. *Compr Psychiatry* 54:568–574.
62. Radloff LS (1991): The use of the Center for Epidemiologic Studies Depression Scale in adolescents and young adults. *J Youth Adolesc* 20:149–166.
63. Henderson D, Poppe AB, Barch DM, Carter CS, Gold JM, Ragland JD, *et al.* (2012): Optimization of a goal maintenance task for use in clinical applications. *Schizophr Bull* 38:104–113.
64. Gold JM, Barch DM, Feuerstahler LM, Carter CS, MacDonald AW 3rd, Ragland JD, *et al.* (2019): Working Memory Impairment across Psychotic disorders. *Schizophr Bull* 45:804–812.
65. Myin-Germeys I, van Os J, Schwartz JE, Stone AA, Delespaul PA (2001): Emotional reactivity to daily life stress in psychosis. *Arch Gen Psychiatry* 58:1137–1144.

66. Bharadhwaj VS, Mubeen S, Sargsyan A, Jose GM, Geissler S, Hofmann-Apitius M, *et al.* (2023): Integrative analysis to identify shared mechanisms between schizophrenia and bipolar disorder and their comorbidities. *Prog Neuropsychopharmacol Biol Psychiatry* 122:110688.
67. Cattarinussi G, Kubera KM, Hirjak D, Wolf RC, Sambataro F (2022): Neural correlates of the risk for schizophrenia and bipolar disorder: A meta-analysis of structural and functional neuroimaging studies. *Biol Psychiatry* 92:375–384.
68. Cheon EJ, Bearden CE, Sun D, Ching CRK, Andreassen OA, Schmaal L, *et al.* (2022): Cross disorder comparisons of brain structure in schizophrenia, bipolar disorder, major depressive disorder, and 22q11.2 deletion syndrome: A review of ENIGMA findings. *Psychiatry Clin Neurosci* 76:140–161.
69. Li Z, Li D, He Y, Wang K, Ma X, Chen X (2022): Cross-disorder analysis of shared genetic components between cortical structures and major psychiatric disorders. *Schizophr Bull* 48:1145–1154.
70. Hindley G, Frei O, Shadrin AA, Cheng W, O'Connell KS, Icick R, *et al.* (2022): Charting the landscape of genetic overlap between mental disorders and related traits beyond genetic correlation. *Am J Psychiatry* 179:833–843.
71. Hill SK, Reilly JL, Keefe RS, Gold JM, Bishop JR, Gershon ES, *et al.* (2013): Neuropsychological impairments in schizophrenia and psychotic bipolar disorder: Findings from the Bipolar-Schizophrenia Network on Intermediate Phenotypes (B-SNIP) study. *Am J Psychiatry* 170:1275–1284.
72. Ventura J, Hellemann GS, Thames AD, Koellner V, Nuechterlein KH (2009): Symptoms as mediators of the relationship between neurocognition and functional outcome in schizophrenia: A meta-analysis. *Schizophr Res* 113:189–199.